

# Determination of the extragalactic background light spectral energy distribution with H.E.S.S.

*Monday, 26 October 2015 17:13 (15 minutes)*

When very high-energy photons (VHE,  $E > 100$  GeV) propagate over cosmological distances, they interact with background light by pair production. Observations of spectral features in the VHE band of extragalactic sources related to this energy-dependent absorption process with the H.E.S.S. array of Cherenkov telescopes allow measuring the spectral energy distribution (SED) of the extragalactic background light (EBL), otherwise very difficult to determine. Preliminary results on the determination of the SED of the EBL will be presented, based on the measurements of the energy spectra of blazars with H.E.S.S. . This model independent approach shows that the shape and overall normalization of the EBL SED is accessible.

**Primary author:** Dr SCHÜSSLER, Fabian (CEA-Saclay / Irfu)

**Co-authors:** Mr SANCHEZ, David (LAPP Annecy, CNRS); Mr LORENTZ, Matthias (IRFU - CEA Saclay); Mr BRUN, Pierre (Irfu - CEA Saclay)

**Presenter:** Dr SCHÜSSLER, Fabian (CEA-Saclay / Irfu)

**Session Classification:** Gamma-Ray Astrophysics

**Track Classification:** Gamma-ray Astrophysics