Type: Oral presentation

High-energy Particle Emission and Cumulative Background from Low-Luminosity Active Galactic Nuclei

Thursday, 29 October 2015 17:55 (15 minutes)

The origin of high-energy neutrinos detected by the IceCube neutrino telescope is a big mystery of highenergy astrophysics. We propose low-luminosity active galactic nuclei (LLAGNs) as a novel source of the high-energy neutrinos. The radiation inefficient accretion flows (RIAFs) are believed to exist in LLAGNs. The Coulomb collisions inside RIAFs is so inefficient that plasmas naturally have high-energy protons through the stochastic particle acceleration. We calculate spectra of escaping neutrinos and cosmic-ray (CR) protons, and find that the RIAFs in LLAGNs can accelerate protons up to $\tilde{}$ PeVs and emit TeV–PeV neutrinos via pp and/or p γ reactions. If $\tilde{}$ 1% of the accretion luminosity is carried away by non-thermal protons, the diffuse neutrino intensity from the RIAFs in LLAGN can be compatible with the observed IceCube data (Kimura, Murase & Toma 2015, ApJ, 806, 159). This result does not contradict either of the diffuse gamma-ray background observed by Fermi or observed diffuse CR flux. This model can be tested by gamma-ray observations around the cores of LLAGNs (e.g., Fujita, Kimura, Murase 2015, PRD, 92, 023001).

Primary author: Dr KIMURA, Shigeo (Tohoku University)

Co-authors: Prof. TOMA, Kenji (Tohoku University); Prof. MURASE, Kohta (The Pennsylvania State University); Prof. FUJITA, Yutaka (Osaka University)

Presenter: Dr KIMURA, Shigeo (Tohoku University)

Session Classification: Neutrinos

Track Classification: Neutrino physics