

# Star-Jet Interactions and Gamma-Ray Flares

Maxim V. Barkov<sup>1</sup>

<sup>1</sup>ABBL RIKEN, Wako, Japan

ICRR University of Tokyo  
26 October 2015

# My collaborators

Felix A. Aharonian

Sergey V. Bogovalov

Valentí Bosch-Ramon

Anton V. Dorodnitsyn

Stanislav R. Kelner

Dmitriy V. Khangulyan


Manel Perucho



# Outline

- 1 VHE short variability and MAIN Ingredients (Jet&Star)
- 2 A low power jet (M87)
- 3 A powerful jet and heavy cloud (3C454.3)
- 4 A few comments about IC310
- 5 Conclusions

# PKS 2155–304 observations



The observed parameters of the PKS 2155–304 flares (H.E.S.S. data)

$$L_\gamma \approx 10^{47} \text{ erg s}^{-1}$$

$$\tau \approx 200 \text{ s}$$


$$L_X \sim 10^{46} \text{ erg s}^{-1}$$

(Aharonian et al 2007)



# What are the Blobs in Powerful Jets?

There are a lot of hypothetical blobs



Internal Shocks, Magnetic Reconnection, Change in Accretion, Instabilities....

# Blobs of external origin

- If blobs have external origin, they can be **very small** as compared to the hydrodynamical scale of the jet....
- External blobs contain **no energy** (as compared to the jet)
- I.e. external blobs must be able to **trigger an intensive interaction**. To be heavy?
- Compact and heavy, i.e **DENSE**: stars, BLR clouds?

Specific realization of such blob formation:

*Jet-Red Giant Interaction Scenario*



# VHE variability in M87





# H.E.S.S., MAGIC, VERITAS observations of M87

Several flashes were observed in 2006, 2008, 2010.

Variability on scales  $t \sim 1\text{ day}$


The flux  $L_\gamma \sim 10^{42}\text{ ergs s}^{-1}$        $E_{\gamma,\max} \simeq 20\text{ TeV}$ .



(Aharonian et al 2006; Abramowski et al. 2011; Aliu et al. 2011)



# Cloud/Star — Jet interaction



(Barkov et al 2010, 2012b)

# Star envelope evolution (Numerical results)

Uniform cloud


(Bosch-Ramon et al 2012)



## p-p interaction



The cloud density can be very high making the  $pp$  interactions to be the most plausible mechanism for the gamma-ray production in the RG-jet interaction scenario: in this case the characteristic cooling time for  $pp$  collisions is

$$t_{pp} \approx \frac{10^{15}}{c_f n_c} = 10^5 n_{c,10}^{-1} c_f^{-1} \text{ s} \quad \chi \equiv E_\gamma/E_p = 0.17 [2 - \exp(-t_\nu/t_{pp})]$$



# VHE light curves and spectra (Numerical model)

$$\xi = 0.5 \text{ and } Q_p(E) \propto E^{-2}$$




(Barkov et al 2012b)



# Fast variability in GeV blazars (3C454.3)

# 3C454.3 observations



The observed parameters of the 3C454.3 flares (*Fermi* data)

$$L_{\gamma} \approx 2 \times 10^{50} \text{ erg s}^{-1}$$




$$\tau_r \approx 4.5 \text{ h}$$

$$L_X \sim 5 \times 10^{47} \text{ erg s}^{-1}$$

(Abdo et al. 2011; Vercellone et al. 2011)





# Dynamical light curve + Radiation spectra: Proton synchrotron and secondary synchrotron




$$t_{\text{acc}}/(2\Gamma_b^2) \approx 5 \text{ h.}$$

# Radiation Model: Geometry




# Radiation Model: Dynamical light curve + cooling time



## A few comments about IC310

# TeV Flare in IC310



The observed parameters of the IC310 TeV flares (MAGIC data)

$$L_\gamma \approx 2 \times 10^{44} \text{ erg s}^{-1}$$

$$\tau_r \approx 4.8 \text{ min}$$

Aleksic et al (2014)




# Black Hole Magnetospheric Model:

## Optimistic Total Energetic Budget

- $\Delta V \lesssim hB_{\text{bh}} \frac{R\Omega_F \sin \theta}{c}, \quad \frac{\Omega_F}{c} \simeq \frac{1}{4r_g}$
- $L_{\gamma,ms} < 4\pi R^2 c \kappa \rho_{\text{GJ}} \Delta V, \quad \rho_{\text{GJ}} = \Omega_F B_{\text{bh}} \sin \theta / (2\pi c)$
- $L_{\gamma,ms} < \frac{1}{8} \kappa B_{\text{bh}}^2 r_g h c \sin^2 \theta$  (2x larger compare to Blandford-Znajek)
- $\dot{m} < 3 \times 10^{-4} M_8^{-1/7}$  the maximum accretion rate compatible with vacuum gap (Levinson&Rieger 2011)
- $B_d = \sqrt{8\pi\beta_m p_g}, \quad h = 10^{13} t_{\text{var},5} \text{ cm}$
- $L_{\gamma,ms} < 10^{43} \beta_m \kappa t_{\text{var},5} M_{\text{BH},8}^{-1/7} \sin^2 \theta \text{ erg s}^{-1}$
- $L_{\gamma,\text{observed}} \approx 2 \times 10^{44} \text{ erg s}^{-1}$

## Black Hole magnetosphere




Aleksic et al (2014)

# Black Hole Magnetospheric Model:

## Optimistic Total Energetic Budget

- $\Delta V \lesssim hB_{\text{bh}} \frac{R\Omega_F \sin \theta}{c}, \quad \frac{\Omega_F}{c} \simeq \frac{1}{4r_g}$
- $L_{\gamma,ms} < 4\pi R^2 c \kappa \rho_{\text{GJ}} \Delta V, \quad \rho_{\text{GJ}} = \Omega_F B_{\text{bh}} \sin \theta / (2\pi c)$
- $L_{\gamma,ms} < \frac{1}{8} \kappa B_{\text{bh}}^2 r_g h c \sin^2 \theta$  (2x larger compare to Blandford-Znajek)
- $\dot{m} < 3 \times 10^{-4} M_8^{-1/7}$  the maximum accretion rate compatible with vacuum gap (Levinson&Rieger 2011)
- $B_d = \sqrt{8\pi\beta_m p_g}, \quad h = 10^{13} t_{\text{var},5} \text{ cm}$
- $L_{\gamma,ms} < 10^{43} \beta_m \kappa t_{\text{var},5} M_{\text{BH},8}^{-1/7} \sin^2 \theta \text{ erg s}^{-1}$
- $L_{\gamma,\text{observed}} \approx 2 \times 10^{44} \text{ erg s}^{-1}$

## Black Hole magnetosphere




Aleksic et al (2014)

# Black Hole Magnetospheric Model:

## Optimistic Total Energetic Budget

- $\Delta V \lesssim hB_{\text{bh}} \frac{R\Omega_F \sin \theta}{c}, \quad \frac{\Omega_F}{c} \simeq \frac{1}{4r_g}$
- $L_{\gamma,ms} < 4\pi R^2 c \kappa \rho_{\text{GJ}} \Delta V, \quad \rho_{\text{GJ}} = \Omega_F B_{\text{bh}} \sin \theta / (2\pi c)$
- $L_{\gamma,ms} < \frac{1}{8} \kappa B_{\text{bh}}^2 r_g h c \sin^2 \theta$  (2x larger compare to Blandford-Znajek)
- $\dot{m} < 3 \times 10^{-4} M_8^{-1/7}$  the maximum accretion rate compatible with vacuum gap (Levinson&Rieger 2011)
- $B_d = \sqrt{8\pi\beta_m p_g}, \quad h = 10^{13} t_{\text{var},5} \text{ cm}$
- $L_{\gamma,ms} < 10^{43} \beta_m \kappa t_{\text{var},5} M_{\text{BH},8}^{-1/7} \sin^2 \theta \text{ erg s}^{-1}$
- $L_{\gamma,\text{observed}} \approx 2 \times 10^{44} \text{ erg s}^{-1}$

## Black Hole magnetosphere



Aleksic et al (2014)

# Black Hole Magnetospheric Model:

## Optimistic Total Energetic Budget

- $\Delta V \lesssim hB_{\text{bh}} \frac{R\Omega_F \sin \theta}{c}$ ,  $\frac{\Omega_F}{c} \simeq \frac{1}{4r_g}$
- $L_{\gamma,ms} < 4\pi R^2 c \kappa \rho_{\text{GJ}} \Delta V$ ,  $\rho_{\text{GJ}} = \Omega_F B_{\text{bh}} \sin \theta / (2\pi c)$
- $L_{\gamma,ms} < \frac{1}{8} \kappa B_{\text{bh}}^2 r_g h c \sin^2 \theta$  (2x larger compare to Blandford-Znajek)
- $\dot{m} < 3 \times 10^{-4} M_8^{-1/7}$  the maximum accretion rate compatible with vacuum gap (Levinson&Rieger 2011)
- $B_d = \sqrt{8\pi\beta_m p_g}$ ,  $h = 10^{13} t_{\text{var},5} \text{ cm}$
- $L_{\gamma,ms} < 10^{43} \beta_m \kappa t_{\text{var},5} M_{\text{BH},8}^{-1/7} \sin^2 \theta \text{ erg s}^{-1}$
- $L_{\gamma,\text{observed}} \approx 2 \times 10^{44} \text{ erg s}^{-1}$
- $L_{\gamma,\text{observed}} \gg L_{\gamma,ms}$

## Black Hole magnetosphere



Aleksic et al (2014)

# Conclusions

- In the case of 3C454.3 the radiation in the GeV energy range can be effectively produced through **proton synchrotron** radiation, Jitter or EIC in the Thompson regime.
- The process can render suitable conditions for energy dissipation and proton acceleration, which could explain the detected day-scale TeV flares in 2010 from M87 via **proton-proton** collisions.
- In IC310 magnetospheric model do **not work**.

Based on:

-  MVB, F.A. Aharonian and V. Bosch-Ramon, (M87); ApJ (2010) 724, 1517
-  MVB, F.A. Aharonian, S.V. Bogovalov, S.R. Kelner and D.V. Khangulyan, (PKS 2155–304 ); ApJ (2012) 749, 119
-  V. Bosch-Ramon, M. Perucho and MVB, (M87); A&A (2012) 539, 69
-  MVB, V. Bosch-Ramon and F.A. Aharonian, (M87); ApJ (2012) 755, 170
-  D.V. Khangulyan, MVB, V. Bosch-Ramon, F.A. Aharonian and A. Dorodnitsyn, (3C454.3) ApJ (2013) 774, 113

# Thank you!!!



# Main Ingredients

## AGN jet

- Relativistic outflow ( $\Gamma_{\text{bulk}} \sim 10 - 100$ , likely depends on the distance)
- Narrow: typically one adopts  $\theta \simeq \Gamma^{-1}$ , i.e.,
- Cross section:

$$\omega \simeq 10^{17} \Gamma_{1.5}^{-1} R_{\text{pc}} \text{cm}$$

## Stars around BH

- Moves with Keplerian velocity:

$$V_* \simeq 600 M_{\text{BH}}^{1/2} R_{\text{pc}}^{-1/2} \text{km/s}$$

- Density (quite uncertain):  $\rho_* \simeq \rho_0 R^{-a}$

Mass injection between  $10^{-2}$  and  $10^{-1}$  pc:

$$\dot{M}_* \simeq 2 \times 10^{-5} \frac{\rho_0 M_{\text{BH},8}^{1/2}}{\Gamma_{1.5}} \int_{0.01}^{0.1} x^{1/2-a} dx [\text{pc}^3 \text{yr}^{-1}],$$



# Probability to get a star to a jet

Murphy et al. 1991

- it was revealed that “ $a$ ” spans a quite broad range depending on the mass accumulated in the central parsec
- It was obtained that  $a = 7/2$  for  $\bar{\rho} = 10^6 M_{\odot} \text{pc}^{-3}$  and  $a = 1/2$  for  $\bar{\rho} = 10^8 M_{\odot} \text{pc}^{-3}$

Mass injection appears to depend very weakly on  $a$


$$\dot{M}_* \simeq 2 \times 10^2 M_{\text{BH},8}^{1/2} M_{\odot} \Gamma_{1.5}^{-1} \text{yr}^{-1}$$

for  $10^{-2} < R_{\text{pc}} < 0.1$

One can expect HUNDREDS of stars entering per year  
which can contain a few Red Giants or young stars per year...



# The Model Solution for the Main Flare







$$D \equiv \frac{L_j r_c^2}{4\theta^2 \Gamma_j^3 z_{0.17} c^3 M_c} \quad L_j \geq 10^{48} \text{ erg s}^{-1}$$


$$M_{\text{BH}} \approx 10^9 M_\odot \quad \delta_b \approx 20$$



# Radiation Model: limitations



# Sketch and Plateau model



$$\dot{M}_* \approx 10^{24} L_{\gamma,49} \xi_{-1}^{-1} \Gamma_{j,1.5}^{-3} \text{ g/s.}$$

The cosmic ray/X-ray excite stellar wind (Basko et al. 1973; Dorodnitsyn et al. 2008),

$$\dot{M} \approx 10^{24} \alpha_{-12} R_{*,2}^{5/2} M_{*,0}^{-1/2} \chi P_{0,6} \text{ g s}^{-1}$$

which providing limitations on the stellar radius

$$R_{*,2} \gtrsim \left( \frac{2 \bar{F}_e M_{0,*}^{1/2}}{\alpha_{-12} \chi} \right)^{2/5}.$$